

Accession #:
Order #:
Reference #:
Patient:
Date of Birth:
Age:
Sex:
Reprinted:
Comment:

Date Collected:
Date Received:
Date of Report:

Telephone:
Fax:



0076 Allergix® Bloodspot IgG4 Food Antibodies 30 Profile

Methodology: ELISA

Negative	Foods to Avoid		
	Mild +1 and +2	Moderate +3 and +4	Severe +5
Aspergillus	Almond	Milk	Egg, Whole
Beef	Mustard	Peanut	
Cantaloupe	Strawberry		
Cashew			
Chicken			
Corn			
Crab			
Garlic			
Lobster			
Oat			
Orange			
Pea, Green			
Pinto Bean			
Pork			
Rice			
Salmon			
Shrimp			
Soybean			
Sunflower			
Tomato			
Tuna			
Turkey			
Walnut			
Wheat			

Responses reflect IgG levels measured by ELISA with standardized food extracts. The assay yields semi-quantitative antibody concentrations for each food. The concentration readings are categorized into four reaction levels (Negative, Mild, Moderate, or Severe) corresponding to semi quantitative responses (0/1, +1, +2, +4 and +5) based on relative absorbance readings. The likelihood of adverse reactions to a given food increases as the response level for that food becomes more positive.